

Spontaneous Oscillations of Collective Molecular Motors

Frank Jülicher and Jacques Prost

Institut Curie, Physicochimie Curie, Section de Recherche, 26 rue d'Ulm, 75231 Paris Cedex 05,

France

(February 1, 2008)

Abstract

We analyze a simple stochastic model to describe motor molecules which cooperate in large groups and present a physical mechanism which can lead to oscillatory motion if the motors are elastically coupled to their environment. Beyond a critical fuel concentration, the non-moving state of the system becomes unstable with respect to a mode with angular frequency ω . We present a perturbative description of the system near the instability and demonstrate that oscillation frequencies are determined by the typical timescales of the motors.

PACS Numbers: 87.10.+e, 05.40.+j

Motor proteins are highly specialized macromolecules which can consume mechanical energy to induce motion and to generate forces. These molecules are involved in active transport processes, cell locomotion and muscle contraction [1,2]. A typical motor molecule specifically attaches to a certain protein filament which serves as a track for its motion. In the presence of fuel, which in the cell is adenosinetriphosphate (ATP), the motor starts moving in a direction defined by the polarity of the track filament. Experimental methods to study the physical properties of motor proteins have been developed in recent years [3–7]. These methods allow to measure forces and velocities of individual motors or small groups of motors.

Normally, at given fuel concentration and temperature, motor molecules will generate a constant average force leading to a constant average velocity [7]. In some cases, however, biological motors are used to drive oscillatory motion. Insect flight muscles e.g. generate oscillating forces in order to move the wings with high frequency. While for some insects (e.g. butterflies) the timing of these oscillations is controlled by a periodic external nerve signal, others (e.g. bees and wasps) generate oscillations within the muscle. The mechanisms which lead to these oscillations are not understood but they seem to be related to the process of force-generation [8].

The purpose of this article is to describe theoretically a physical mechanism by which motor collections acting on a spring can give rise to oscillatory motion. Oscillations occur as the result of cooperation of many motors while individual motors only generate time-independent velocities.

The microscopic mechanisms which are responsible for the force generation of molecular motors are not understood. In order to identify possible physical mechanisms which allow such a motor molecule to convert the chemical energy of a fuel to motion and mechanical work, simple physical models have been suggested [9–16]. One idea [10] is to simplify the internal degrees of freedom of the motor to two different states of a particles which moves along a one-dimensional coordinate x . The interaction of the particle with the track is described by periodic potentials which reflect the periodic structure along the surface of the

track. Within such a two-state model motion is induced if the potentials are asymmetric with respect to $x \rightarrow -x$ (which reflects the polarity of the track) and if detailed balance of the transitions between states is broken [10].

New phenomena occur if the collective motion of many particles is considered [17–19]. In a simple model for collective motion, particles are rigidly coupled to a common structure which they set in motion collectively [18]. As a result of cooperation, motion can in this case even occur in a symmetric system via spontaneous symmetry breaking. Instabilities and dynamical phase transitions also occur in asymmetric systems revealing the rich behaviour of many-motor systems. At these transitions, a steady state becomes unstable and the system chooses a new steady state. These transitions correspond therefore to dynamical instabilities at angular frequency $\omega = 0$.

In the following, we demonstrate that many coupled motors also can induce instabilities with $\omega \neq 0$ towards a state with periodically varying velocity. Such instabilities occur if the system is elastically coupled to its environment. This implies, that the complete system is connected to its environment via a spring attached to the common backbone, see Fig. 1 [20]. A single particle reaches in this case a steady state with zero velocity and a position at which the average force generated by the motor is balanced by the elastic force of the spring [5].

We are interested in the behavior of many motors coupled together. Let us first review the most simple two-state model which we are going to discuss [10,18]. Motors are described by particles which move in two periodic potentials $W_\sigma(x)$ with period l . Transitions between states $\sigma = 1, 2$ occur with rates $\omega_1(x)$ and $\omega_2(x)$ which also are l -periodic. The excitation amplitude can in general be written as [10] $\omega_1(x) = \omega_2(x)[\exp[(W_1 - W_2)/T] + \Omega\theta(x)]$, where T denotes temperature measured in units of the Boltzmann constant. We assume that the periodic function $\theta(x)$ is given. In principle it can be calculated from a specific model for the reaction kinetics of the fuel coupled to the particle [21]. The amplitude Ω is related to the difference of chemical potentials $\Delta\mu \equiv \mu_{ATP} - \mu_{ADP} - \mu_P$ of the fuel and its products which is the chemical driving force of the motors. Here, we assume that the fuel is ATP which

is hydrolyzed to adenosinediphosphate (ADP) and phosphate (P): $ATP \rightleftharpoons ADP + P$. For $\Delta\mu = 0$, $\Omega = 0$ and the rates ω_1 and ω_2 obey detailed balance $\omega_1/\omega_2 = \exp[(W_1 - W_2)/T]$. As soon as $\Delta\mu \neq 0$, Ω becomes nonzero. For small $\Delta\mu/T$, $\Omega \sim \Delta\mu$; for large $\Delta\mu/T$, $\Omega \sim C_{ATP}$ is proportional to the fuel concentration.

We now study the behavior of many particles which are rigidly connected to a common backbone, see Fig. 1. Particles have a fixed spacing q which for simplicity is assumed to be incommensurate with the period l , i.e. l/q is irrational. In the limit of an infinite system one can then introduce densities $P_\sigma(\xi)$ with $\sigma = 1, 2$ which give the probability to find a particle at position $\xi = x \bmod l$ relative to the beginning of the potential period in state σ . These densities are not independent but obey $P_1(\xi) + P_2(\xi) = 1/l$ and $\int_0^l d\xi (P_1 + P_2) = 1$. The equations of motion for this system read [18]

$$\begin{aligned} \partial_t P_1 + v \partial_\xi P_1 &= -\omega_1 P_1 + \omega_2 P_2 \\ \partial_t P_2 + v \partial_\xi P_2 &= \omega_1 P_1 - \omega_2 P_2 \end{aligned} \tag{1}$$

$$f_{ext} = \lambda v + KX + \int_0^l d\xi P_1 \partial_\xi (W_1 - W_2) \quad . \tag{2}$$

Eq. (1) describes the dynamics of the density $P_1(\xi)$ resultig from motion of the backbone with velocity $v = \partial_t X$ and the transitions between the states. Eq. (2) is a force balance: The externally applied force per particle $f_{ext}(t)$ is balanced by viscous drag with damping coefficient λ , the average force exerted by the potentials and an additional elastic force KX of a spring of length X with elastic modulus KN where N is the number of particles. Note, that Eqns. (1) and (2) only depend on the difference $W_1 - W_2$ of the two potentials and we can choose W_2 to be constant without loss of generality.

As a consequence of the spring action, a nonmoving solution to Eqns. (1) and (2) exists for $f_{ext} = 0$, $v = 0$, $P_1 = R \equiv \omega_2/(\alpha l)$, where $\alpha(\xi) \equiv \omega_1(\xi) + \omega_2(\xi)$ and $X = X_0 \equiv -\int_0^l d\xi R \partial_\xi (W_1 - W_2)/K$. We first study the linear stability of this solution and determine the instability threshold with respect to oscillations. With the ansatz $P_1(\xi, t) = R(\xi) + p(\xi) \exp(st)$, $v(t) = u \exp(st)$ and $X(t) = X_0 + u \exp(st)/s$, one finds using Eq. (1) and (2) to linear order in p and the velocity amplitude u :

$$p(\xi) = -u \frac{\partial_\xi R}{s + \alpha(\xi)} . \quad (3)$$

The possible values of the complex eigenvalue $s = -\tau + i\omega$ are determined by

$$\lambda + \frac{K}{s} = \int_0^l d\xi \frac{\partial_\xi R \partial_\xi (W_1 - W_2)}{s + \alpha(\xi)} . \quad (4)$$

The non-moving state is unstable if $\tau < 0$. The instability occurs for $\tau = 0$, where the real part of s vanishes. This happens at a threshold value $\Omega = \Omega_c(K)$ for which $\alpha(\xi) = \alpha_c(\xi)$.

This critical value together with the frequency ω_c at the instability is determined by

$$\lambda = \int_0^l d\xi \frac{\alpha_c}{\alpha_c^2 + \omega_c^2} \partial_\xi R \partial_\xi (W_1 - W_2) \quad (5)$$

$$K = \int_0^l d\xi \frac{\omega_c^2}{\alpha_c^2 + \omega_c^2} \partial_\xi R \partial_\xi (W_1 - W_2) . \quad (6)$$

In the limit $K = 0$ where no elastic element is present, it follows from Eq. (6) that $\omega_c = 0$. Eq. (5) now determines $\Omega_c(0)$ which is the condition for the instability of the non-moving state as previously described in Ref. [18]. In the case of a symmetric system, this instability leads to spontaneous motion via a symmetry-breaking transition. For nonzero K the instability occurs for $\Omega_c(K) = \Omega_c(0) + \delta\Omega_c$ with $\delta\Omega_c \sim K$ for small K and the system starts to oscillate with finite angular frequency $\omega_c \sim K^{1/2}$. Note, that beyond a maximal value $K > K_{max}$, the resting state is stable for any value of Ω .

Fig. 2 (a) and (b) show examples for these oscillations which have been calculated numerically for constant deexcitation rate ω_2 , $\lambda\omega_2 l^2/U = 0.1$ and piecewise linear and symmetric potentials $W_\sigma(\xi)$, see Fig. 1 (a) with $a = l/2$. In this case $K_{max} l^2/U \simeq 2.8$. The function $\theta(\xi)$ is chosen as shown in Fig. 1 (a) with $d/l = 0.1$. Fig. 2 (a) displays the position X versus time t for an excitation level $\Omega = 0.1$ slightly above threshold and elastic modulus $Kl^2/U = 0.2$. After an initial relaxation period, motion is almost sinusoidal. An example for small elastic modulus $Kl^2/U = 0.002$ and same Ω is shown in Fig. 2 (b). The periodic motion in this case shows cusp-like maxima which correspond to sudden changes of the velocity. These cusps are related to discontinuities of the velocity as a function of external force for $K = 0$ [18].

We now give an analytical description of oscillations in the vicinity of the instability. Anticipating, that the motion is periodic with period $t_P \equiv 2\pi/\omega$, we can write $P_1(\xi, t) = \sum_{k=-\infty}^{\infty} P_1(\xi, k)e^{ik\omega t}$, $v(t) = \sum_{k=-\infty}^{\infty} v_k e^{ik\omega t}$, and $f_{ext}(t) = \sum_{k=-\infty}^{\infty} f_k e^{ik\omega t}$, which defines the Fourier coefficients $P_1(\xi, k)$, v_k and f_k . Using this representation, one can derive the nonlinear relation between velocity and external force

$$f_k = F_{kl}^{(1)}v_l + F_{klm}^{(2)}v_lv_m + F_{klmn}^{(3)}v_lv_mv_n + O(v^4) \quad . \quad (7)$$

The coefficients $F_{k,k_1,\dots,k_n}^{(n)}$ can be calculated by first rewriting Eq. (1) as

$$P_1(\xi, k) = \delta_{k,0}R(\xi) - \sum_{lm} \frac{\delta_{k,l+m}}{\alpha + i\omega k} v_l \partial_\xi P_1(\xi, m) \quad . \quad (8)$$

Inserting the ansatz

$$P_1(\xi, k) = R\delta_{k,0} + P_{kl}^{(1)}(\xi)v_l + P_{klm}^{(2)}(\xi)v_lv_m + O(v^3) \quad , \quad (9)$$

into Eq. (8), one obtains a recursion relation for the functions $P_{k,k_1,\dots,k_n}^{(n)}$:

$$P^{(n)}(\xi)_{k,k_1,\dots,k_n} = - \sum_l \frac{\delta_{k,k_n+l}}{\alpha + i\omega k} \partial_\xi P_{l,k_1,\dots,k_{n-1}}^{(n-1)} \quad (10)$$

Using Eq. (2), one finds

$$F_{kl}^{(1)} \equiv \delta_{kl} \left(\lambda + \frac{K}{i\omega k} - \int_0^l d\xi \frac{\partial_\xi R \partial_\xi (W_1 - W_2)}{\alpha + i\omega k} \right) \quad . \quad (11)$$

The linear response function of the system is given by the inverse of $F_{kl}^{(1)}$. Therefore the instability condition Eq. (4) corresponds to $F_{kl}^{(1)} = 0$. For $n > 1$,

$$F_{k,k_1,\dots,k_n}^{(n)} \equiv \int_0^l d\xi P^{(n)}(\xi)_{k,k_1,\dots,k_n} \partial_\xi (W_1 - W_2) \quad . \quad (12)$$

Note, that the coefficients $F_{k,k_1,\dots,k_n}^{(n)}$ are nonzero only if $k = k_1 + \dots + k_n$. This results from time translational symmetry [22]. If no elastic coupling to the environment is present, i.e. $K = 0$, one recovers for constant external force f_0 the steady states which have been described previously [18]: $f_{ext} = f_0(v_0)$; $v_n = 0$ for $n \neq 0$. As soon as $K \neq 0$, v_0 must vanish and a constant external force f_0 only changes the average position X_0 .

Spontaneous oscillations are solutions to Eq. (7) for $f_n = 0$. The dominant terms near the instability of v_1 are given by

$$0 = F_{11}^{(1)}v_1 + G^{(2)}v_{-1}v_2 + G^{(3)}v_1^2v_{-1} \quad (13)$$

$$0 = F_{22}^{(1)}v_2 + F_{211}^{(2)}v_1^2 \quad , \quad (14)$$

where $G^{(2)} \equiv F_{1,2,-1}^{(2)} + F_{1,-1,2}^{(2)}$ and $G^{(3)} \equiv F_{1,1,1,-1}^{(3)} + F_{1,1,-1,1}^{(3)} + F_{1,-1,1,1}^{(3)}$. As soon as v_1 and v_2 are determined, higher orders v_n can be obtained recursively using Eq. (7). From Eq. (14) one finds that $v_2 \sim v_1^2$. Inserting this value in Eq. (13), one obtains

$$0 = F_{11}^{(1)}v_1 + \tilde{G}^{(3)}v_1^2v_{-1} \quad , \quad (15)$$

with an effective third order coefficient $\tilde{G}^{(3)} \equiv G^{(3)} - F_{211}^{(2)}G^{(2)}/F_{22}^{(1)}$. One solution to Eq. (15) is always $v_1 = 0$. The remaining solutions are described by

$$|v_1|^2 = -F_{11}^{(1)}/\tilde{G}^{(3)} \quad . \quad (16)$$

Since the amplitude $|v_1|^2$ is a real number, solutions to Eq. (16) exist only if the right hand side of Eq. (16) is real and positive. In general, however, $F_{11}^{(1)}$ and $\tilde{G}^{(3)}$ are complex numbers. The requirement to find real solutions to Eq. (16) fixes the oscillation frequency. This can be seen as follows: If the system is at its instability threshold $\Omega = \Omega_c(K)$, $F_{11}^{(1)} = 0$ for $\omega = \omega_c$, see Eq. (4). In the (Ω, ω) -plane, a line exists along which $F_{11}^{(1)}/\tilde{G}^{(3)}$ is real. This line can be parametrized by a function $\omega = \omega_s(\Omega)$ which at the instability threshold obeys $\omega_c = \omega_s(\Omega_c)$. For $\Omega > \Omega_c$, $-F_{11}^{(1)}/\tilde{G}^{(3)}$ is real and positive along this line thus allowing for a solution to Eq. (16). For $\Omega < \Omega_c$, $v_1 = 0$ is the only solution. Fig. 2 (c) shows the dependence of the selected frequency ω_s as a function of the excitation amplitude. For $\Omega < \Omega_c(K)$, $\omega_s = 0$. At the instability, a finite frequency is selected which increases for increasing Ω .

Note, that this example is developed for a symmetric system for which an additional restriction holds: Eq. (7) is now invariant with respect to the transformation $v_n \rightarrow -v_n$, $f_n \rightarrow -f_n$. As a result, all even expansion coefficients in Eq. (7) vanish, $F^{(2)} = F^{(4)} = \dots = 0$. In particular, $G^{(2)}$ vanishes and $\tilde{G}^{(3)} = G^{(3)}$. Furthermore, all terms which couple v_1 to modes

v_n with even n vanish and $v_2 = v_4 = \dots = 0$. As a result, the periodic motion of a symmetric system has the symmetry property $v(t + t_P/2) = -v(t)$.

The parameters used in Fig. 2 correspond e.g. to the choice $\lambda \simeq 10^{-8} kg/s$, $\omega_2 \simeq 10^3 s^{-1}$, $l \simeq 10^{-8} m$, $U \simeq 10 k_B T$. Oscillation frequencies in this case vary between $\omega_s = 0$ for $K = 0$ and $\omega_s \simeq 5\omega_2$ for $K \simeq K_{max} \simeq 10^{-3} N/m$ per motor [23]. This shows that using typical time scales of biological motors the mechanism described here generates frequencies up to the kHz range.

In summary, we have shown using a simple two-state model for force-generation of molecular motors that oscillatory motion can occur as a collective effect. Motors which individually only generate constant forces, show oscillating behavior if they are coupled in large numbers and work against an elastic spring. Beyond a critical amplitude of excitations which drive the system, a nonmoving state becomes unstable with respect to periodic motion. This argument suggests that oscillations are a possible mode of motion of biological motor proteins and gives an example of how some insect flight muscles could generate oscillations via the mechanism of force generation.

We thank F. Gittes, A. Ott, L. Peliti, D. Riveline and W. Zimmermann for useful discussions and A. Ajdari for a critical reading of the manuscript. We are also very grateful to F. Amblard, M. Bornens, J. Käs and D. Louvard for information concerning spontaneous oscillations in biological systems. F.J. acknowledges a Marie Curie Fellowship of the European Community; part of this work was done at the Aspen center for physics.

REFERENCES

- [1] B. Alberts, D. Bray, J. Lewis, M. Raff, K. Roberts and J.D. Watson, The molecular biology of the cell, (Garland, New York, 1994).
- [2] H.E. Huxley, Science **164**, 1365 (1969).
- [3] J.A. Spudich, Nature **348**, 284 (1990).
- [4] A. Ishijima, T. Doi, K. Sakurada and T. Yanagida, Nature **352** 301 (1991).
- [5] K. Svoboda, C.F. Schmidt, B.J. Schnapp and S.M. Block, Nature **365**, 721 (1993).
- [6] A.J. Hunt, F. Gittes and J. Howard, Biophys. J. **67**, 766 (1994).
- [7] D.A. Winkelmann, L. Bourdieu, A. Ott, F. Kinose, A. Libchaber, Biophys. J. **68**, 2444 (1995); L. Bourdieu, M.O. Magnasco, D.A. Winkelman and A. Libchaber, Phys. Rev. E **52**, 6573 (1996).
- [8] J.W.S. Pringle, in *Insect Flight Muscle*, R.T. Tregear, Ed. , North-Holland, Amsterdam, 177 (1977).
- [9] A. Ajdari and J. Prost, C.R. Acad. Sci. Paris, t.315, Série II, p. 1635 (1992).
- [10] J. Prost, J.-F. Chauwin, L. Peliti and A. Ajdari, Phys. Rev. Lett. **72**, 2652 (1994); J.-F. Chauwin, A. Ajdari and J. Prost, Europhys. Lett. **27**, 421 (1994).
- [11] C.S. Peskin, G.B. Ermentrout, G. Oster, in Cell Mechanics and Cellular Engineering, ed. by V. Mow et. al., Springer, New York (1994).
- [12] R.D. Astumian and M. Bier, Phys. Rev. Lett. **72**, 1766 (1994).
- [13] M.O. Magnasco, Phys. Rev. Lett. **71**, 1477 (1993); M.O. Magnasco, Phys. Rev. Lett. **72**, 2656 (1994).
- [14] C.R. Doering, Nuovo Cimento **17**, 685 (1995).
- [15] I. Derényi and T. Vicsek, Proc. Nat. Acad. Sci. **93**, 6775 (1996).

- [16] H.X. Zhou and Y. Chen, Phys. Rev. Lett **77**, 194 (1996).
- [17] S. Leibler and D. Huse, J. Cell Biol. **121**, 1357 (1993).
- [18] F. Jülicher and J. Prost, Phys. Rev. Lett., **75**, 2618-2621 (1995).
- [19] I. Derényi and T. Vicsek, Phys. Rev. Lett. **75**, 374 (1995); I. Derenyi, A. Ajdari, Phys. Rev. E, **54**, 5-8 (1996).
- [20] Note, that at the microscopic length scales of the motors inertia is negligible and motion is overdamped. Our approach is therefore in contrast to models where inertia terms are needed to obtain oscillations, see e.g. S. Scilia and D.A. Smith, Math. Biosciences **106**, 159 (1991).
- [21] F. Jülicher, A.Ajdari and J. Prost, in preparation.
- [22] This symmetry corresponds to the invariance of Eq. (7) under the transformations $v_n \rightarrow v_n e^{in\phi}$ and $f_n \rightarrow f_n e^{in\phi}$ for arbitrary ϕ .
- [23] Here, we have assumed the limit of large Ω corresponding to saturated ATP concentration. For a system of 10^3 motors, K_{max} corresponds to an elastic modulus of $1N/m$.

FIGURES

FIG. 1. (a) Periodic potentials W_1 and W_2 used to calculate the behavior of the system. The Function $\theta(x)$ is chosen to be nonzero in the vicinity of the potential minimum. (b) Many particles coupled rigidly to a backbone which is connected with its environment via a spring K . The particle spacing is denoted q .

FIG. 2. (a) Position X versus time t for a symmetric system with $d/l = 0.1$, $\Omega = 0.1$, $\lambda\omega_2 l^2/U = 0.1$ and $Kl^2/U = 0.2$ (see Fig. 1 (a)). (b) Position versus time for same system but $Kl^2/U = 0.002$. (c) Selected frequency ω_s as a function of the excitation amplitude Ω for the case $\Omega = 0.1$, $Kl^2/U = 0.2$. The critical excitation amplitude is $\Omega_c \simeq 0.08$.





Fig. 1 (Julicher & Prost)




Fig. 2 (a) (Julicher & Prost)




Fig. 2 (b) (Julicher & Prost)




Fig. 2 (c) (Julicher & Prost)